



The Influence Educational Status on Genital Tract Infections

Corresponding Author:

Dr. Emeka E Ojiyi,
FWACS, FMCOG, FICS, Obstetrics and Gynaecology, Imo State University Teaching Hospital Orlu - Nigeria

Submitting Author:

Dr. Emeka E Ojiyi,
FWACS, FMCOG, FICS, Obstetrics and Gynaecology, Imo State University Teaching Hospital Orlu - Nigeria

Article ID: WMC002558

Article Type: Research articles

Submitted on: 02-Dec-2011, 05:32:08 PM GMT **Published on:** 03-Dec-2011, 08:05:24 PM GMT

Article URL: http://www.webmedcentral.com/article_view/2558

Subject Categories: CYTOLOGY

Keywords: Cervical cytology, Genital tract infections, Educational status

How to cite the article: Ojiyi E E, Dike E I, Anolue F C, Okeudo C, Uzoma O I, Uzoma M J, Agu P O, Anozie U M, Nggada H. The Influence Educational Status on Genital Tract Infections. WebmedCentral CYTOLOGY 2011;2(12):WMC002558

Source(s) of Funding:

The article was funded by the contribution of the authors involved. No grant was received from any external individual or organisation.

Competing Interests:

The authors declare that we do not have any competing interests.

The Influence Educational Status on Genital Tract Infections

Author(s): Ojiyi E E, Dike E I, Anolue F C, Okeudo C, Uzoma O I, Uzoma M J, Agu P O, Anozie U M, Nggada H

Abstract

Background: Genital tract infections such as Gardnerella vaginalis vaginosis, trichomoniasis and candidiasis have continued to assume importance because of the particular unique features of their main causative organisms- Gardnerella vaginalis, Trichomonas vaginalis and Candida albicans, and myriads of clinical and pathological changes in affected persons.

Aim: To study the association between educational status and genital tract infections

Methods: The subjects were 450 randomly selected sexually active women attending antenatal, postnatal, gynecology and family planning clinics in the Department of Obstetrics and Gynecology of the University of Maiduguri Teaching Hospital from April 2001 to May 2002. The Pap smear of these patients were examined microscopically for the evidence of Candida albicans, Gardnerella vaginalis and Trichomonas vaginalis. A questionnaire assessing the educational status of the patients was administered.

Results: Sixty-six (66) of the 450 patients studied had specific infections giving a prevalence rate of 14.7%. Candida albicans, Gardnerella vaginalis, Trichomonas vaginalis and Candida albicans in conjunction with Gardnerella vaginalis contributed 7.1%, 5.15%, 2.4% and 1.8% respectively to this overall prevalence. Their ages ranged between 15 and 64 years, with a mean of 26 ± 3 years.

Conclusion: Those with post secondary educational qualification are at increased risk of acquiring genital tract infections and should be the target in sporadic or organized sexually transmitted infections screening without discriminating other sexually active women.

Key words: Cervical cytology, genital tract infections, educational status.

Introduction

Genital tract infections such as Gardnerella vaginalis vaginosis, trichomoniasis and candidiasis have continued to assume immense importance not only because of the particular unique features of their main causative organisms- Gardnerella vaginalis, Trichomonas vaginalis and Candida albicans, but also

the fact that the organisms can elicit myriads of clinical and pathological changes in different body tissues, organs and systems. [1] They have been implicated in neonatal meningitis, urinary tract infections, pyogenic liver abscess and following prostatectomy. [2-5] Cervical cytology has not only served the purpose of screening for cervical intra-epithelial (CIN) lesions but has been useful in the diagnosis of genital tract infections such as Gardnerella vaginalis, Trichomonas vaginalis and Candida albicans. The Papanicolaou smear for cervical cytology is cost effective, acceptable to most patients and adaptable to wide spread screenings. [7] In low resource areas where facilities might not be easily available to detect the above mentioned genital tract infections, cervical cytology has been found to very helpful. [6] It is also specific enough to detect cytological changes pathognomonic of human papilloma virus, which ordinarily is difficult to isolate under normal laboratory conditions, except by the use of special techniques like DNA Hybridization, PCR amplification or viral culture. [7-9]

Many studies have tried to show some kind of association between occupation and the risk of developing genital tract infections. [6, 10] Such socio-demographic factors may be useful in risk scoring systems. This is important because risk scoring systems have the potential for assisting the targeting of screening resources as broad risk targeting of all sexually active women is not a viable option for developing countries due to paucity of both human and financial resources. Even in the industrialized nations Of the West, the need for more precise targeting of high risk groups in order to improve the efficiency of sexually transmitted infections screening and conserve funds has become a major issue. [11]

This study looks at the association between educational status and genital tract infections.

Materials/Methods

The subjects were 450 randomly selected sexually active women attending various clinics in the department of Obstetrics and Gynaecology of the University of Maiduguri Teaching Hospital.

These included the antenatal, postnatal, gynaecology

and family planning clinics. They were recruited after consenting to participate and a formal approval had been given by the institution's ethics and research committee. The recruitment continued until a sample size of 450 was reached. This was calculated using the Epi Info version 6 programme for population or descriptive study using simple random sampling. It was based on a population of 4,342 patients/ clients attending the recruiting clinics from April 2001 to May 2002. The purpose, value and nature of the procedure was explained to each prospective patient and her consent obtained. All the consenting patients had their pap smears taken using a moistened unlubricated Cusco's bivalve speculum and an Ayre's wooden spatula after a questionnaire containing occupation of the woman had been filled. The smears were immediately transported to the histopathology laboratory immersed in 95% ethanol for preparation, staining and reading. The smears were examined microscopically by a pathologist at the magnifications of 4, 10 and 100.

The WHO Epi Info statistical programme was used to compute and analyze the results. These included frequency distribution and tests of significance using chi-square (χ^2). A P value of

Inclusion criteria: All sexually active women attending the above clinics and who consented to participate in the programme were included until the sample size was reached.

Exclusion criteria: Women who declined to consent were exempted, so were those who had never been sexually exposed. In addition, those with obvious cervical lesions, vaginal discharge and those who were menstruating were excluded from the study.

Results

Table 1 shows the cytology results of the Papanicolaou smears. Specific infections were identified in 66 of the 450 patients studied giving an overall prevalence rate of 14.7%. The contributions each of *Candida albicans*, *Gardnerella vaginalis* and *Trichomonas vaginalis* to this overall prevalence rates is depicted in Table 2, with *Candida albicans* constituting the highest of prevalence of 7.1% of all infections. Table 3 shows the distribution of the patients by educational status. It indicates that the prevalence of the specific genital tract infections increased with increasing educational status, being lowest among those with no formal and primary education and highest in the patients with postsecondary qualifications. This difference was statistically significant ($P = 0.034$, X^2 for linear trend=

0.90).

Discussion

The prevalence of specific genital infections in this study which was 14.7% is similar to the 13.7% reported by Konje et al [6] from Ibadan and Ngokerre and Ofordile [12] from Enugu. The prevalence of *Gardnerella vaginalis* the major aetiological agent of bacterial vaginosis, in this study of 5.1% is much lower than the 40.8% reported by Adinma et al [1] from Nnewi, 42% by Oji [13] from Lagos, and 10% by Chowdry et al [14] in India. Variations in incidence are invariably related to the characteristics of the population studied, the period of study, the technique used in isolating the organisms and the presence or absence of symptoms of the infections among the patients studied. The low prevalence of genital infections in this study compared to those of Adinma et al [1], Oji [13] and Abudu [15] et al is probably due to the fact that virtually all the patients (87%) were asymptomatic. This study shows that the highest incidence of sexually transmitted genital infections (e.g. *T. vaginalis*) increased with increasing levels of educational qualification and is indeed significantly higher among those whom never had any form of formal education. The underlying factor might be multiple sexual partners as those who engage in sexual promiscuity do so for economic reasons [11]. Patients in secondary and tertiary institutions by reason of their age and the influence of reduced or absent of parental control are more likely to be more sexually promiscuous, especially in these days of worsening economic difficulty and decaying societal core values. Some of these patients even train themselves in school. These associations may allow for targeting of a high risk group in sporadic sexually transmitted infections (STI) screening programmes as is the practice in most developing countries such as ours or in an organized systematic STI screening programme without discriminating other sexually active women. There is also need for an increased use of this cost effective and acceptable diagnostic tool in the diagnosis of genital tract infections, especially in centers with organized cervical cytology screening programmes and in low resource areas where there might be shortage of manpower and facilities might not be adequate since it is essentially an office procedure.

References

1. Adinma JI, Okwoli RN, Agbai AO, Unaeze NC.

- Gardnerella vaginalis vaginosis in Nigerian Igbo women. *Trop J Obstet Gynaecol* 2000; 17: 21-23.
2. Berardi- Grassias L, Roy O, Berardi JC, Furioli J. Neonatal meningitis due to Gardnerella vaginalis. *Eur J Clin Microbiol Infect Dis* 1988; 7: 406-407.
 3. Moller BR, Kristianasen FV, Theresen P, Frost L, Morgensen SC. Sterility of the uterine cavity. *Acta Obstet Gynaecol Scand* 1995; 74: 215-219.
 4. Lam HM, Birth DF, Fairley KF. Prevalence of Gardnerella vaginalis in the urinary tract. *J Clin Microbiol* 1988; 26: 1130-1133.
 5. Ezzel JH, Many WJ Jr. Gardnerella vaginalis: an unusual cause of pyogenic liver abscess. *Am J Gastroenterol* 1988; 83: 1409-1411.
 6. Konje JC, Otolarin EO, Ogunniyi JO, Obisesan KA, Ladipo OA. The prevalence of Gardnerella vaginalis, Trichomonas vaginalis and Candida albicans in the cytology clinic in Ibadan, Nigeria. *Afr J Med Med Sci* 1991; 20:29-34.
 7. Guidozi F. Screening for cervical cancer. *Obstet Gynaecol Surv* 1996; 51:247-252.
 8. Singer A, Jekins D. Viruses and cervical cancer. *BMJ* 1991; 302:251-252.
 9. Odujinrin OM, Oyediran MA. Pap smears of contracepting women in Lagos, Nigeria. *Niger Med J* 1996; 30:15-19.
 10. Bro F. Vaginal microbial flora in women with and without discharge registered in general practice. *Danish Med Bull* 1989; 36: 483-485.
 11. Garner HL, Damper TK, Dikes CD. The prevalence of vaginitis. *Am J Obstet Gynaecol* 1957; 73: 1080-1087.
 12. Ngokerre AA, Ofordile PM. Cytological evaluation of cervical smears in the University of Nigeria Teaching Hospital, Enugu and environs: a 5-year study. *Orient J Med* 1996; 8:49-52.
 13. Oji GO. Gardnerella vaginalis in non-specific vaginitis. Its occurrence, identification and antimicrobial susceptibility. M. Sc. Thesis. University of Lagos, 1983.
 14. Chowdhury A, Bhujwala RA, Shrinivas. Gardnerella vaginalis: Isolation and identification. *Indian J Pathol Microbiol* 1990; 33:151-156.
 15. Abudu O, Odugbemi TO, Agboola A. Corynebacterium vaginalis and vaginitis. *West Afr Med* 1982; 1:17-19.
 16. Rotkin ID. Epidemiology of cancer of the cervix III. Sexual characteristics of a cervical screening population. *Am J Pub Health* 1967; 57: 815-829.
 17. Martin CE. Epidemiology of cancer of the cervix II. Marital and coital factors in cervical cancer. *Am J Pub Health* 1967; 57: 803-814.

Illustrations

Illustration 1

Table1. Result of Pap smear in 450 subjects

Class of Pap smear	No	% age
Normal	20	45.6
Specific Infections	66	14.7
Non specific Inflammation	58	12.9
CIN	73	16.2
HPV changes	48	10.7
Total	450	100

Illustration 2

Table2. Percentage distribution of the organisms

Organism	No	%
Candida albicans	3	7.1
Gardnerella vaginalis	23	5.1
Trichomonas vaginalis	11	2.4
Candida albicans+ Gardnerella vaginalis	8	1.8
Overall prevalence	14.7%	N=450

Illustration 3

Table3. Prevalence of infections by educational status

Educational level	Negative	Positive	Total
None	10	102	112
Primary	8	57	65
Secondary	18	172	190
Post- secondary	12	71	83
Total	48	402	450
X² for linear teand= 0.90, df=3			
P=0.034.			

Disclaimer

This article has been downloaded from WebmedCentral. With our unique author driven post publication peer review, contents posted on this web portal do not undergo any prepublication peer or editorial review. It is completely the responsibility of the authors to ensure not only scientific and ethical standards of the manuscript but also its grammatical accuracy. Authors must ensure that they obtain all the necessary permissions before submitting any information that requires obtaining a consent or approval from a third party. Authors should also ensure not to submit any information which they do not have the copyright of or of which they have transferred the copyrights to a third party.

Contents on WebmedCentral are purely for biomedical researchers and scientists. They are not meant to cater to the needs of an individual patient. The web portal or any content(s) therein is neither designed to support, nor replace, the relationship that exists between a patient/site visitor and his/her physician. Your use of the WebmedCentral site and its contents is entirely at your own risk. We do not take any responsibility for any harm that you may suffer or inflict on a third person by following the contents of this website.